


AD-A203 931

DTIC FILE COPY

①

USAARL Report No. 89-1



**Anomalous Retinal Correspondence
(Reprint)**

By

Morris R. Lattimore, Jr.

Sensory Research Division

October 1988

**DTIC
ELECTE
S 24 JAN 1989 D
CE**

89 1 23 146

Approved for public release; distribution unlimited.

**United States Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-5292**

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U.S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this document when it is no longer needed. Do not return it to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Reviewed:



BRUCE C. LEIBRECHT, Ph.D.
LTC, MS
Director, Sensory Research
Division

Released for publication:



J. D. LAMOTHE, Ph.D.
COL, MS
Chairman, Scientific
Review Committee



DAVID H. KARNEY
Colonel, MC
Commanding

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 89-1			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Aeromedical Research Laboratory		6b. OFFICE SYMBOL (If applicable) SGRD-UAS-VS	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Development Command		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 577 Fort Rucker, AL 36362-5292			7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21701-5012		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
11. TITLE (Include Security Classification) Anomalous Retinal Correspondence					
12. PERSONAL AUTHOR(S) Lattimore, Morris R.					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1988 October	
15. PAGE COUNT 11					
16. SUPPLEMENTARY NOTATION This work was done at the University of Houston and this report is a reprint of a publication in Southern Journal of Optometry, Volume VI, Number 3, Summer, 1988					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) anomalous retinal correspondence, strabismus, diagnosis, therapy, theories		
FIELD	GROUP	SUB-GROUP			
06	04				
06	05				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This paper presents an overview of anomalous retinal correspondence in strabismus. Definitions, certain testing techniques, and a review of underlying theory are outlined. It is concluded that ARC is not well understood and represents an area still open for investigation. <i>Reprints 1/89</i>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Scientific Information Center			22b. TELEPHONE (Include Area Code) (205) 255-6907		22c. OFFICE SYMBOL SGRD-UAX-SI

ANOMALOUS RETINAL CORRESPONDENCE

MORRIS R. LATTIMORE, JR., O.D.
U.S. Army Aeromedical Research Laboratory

Accession For

NTIS GRA&I

DTIC TAB

Unannounced

Justification

Distribution/

Availability Codes

Dist

Avail and/or
Special

A-120

Abstract. This paper presents an overview of anomalous retinal correspondence in strabismus. Definitions, certain testing techniques, and a review of underlying theory are outlined. It is concluded that ARC is not well understood and represents an area still open for investigation.

Keywords: Diagnosis
Medicine, Therapy, reprints. (AW)

DTIC

COPY
INSPECTED
A

INTRODUCTION

Anomalous correspondence has been a topic of considerable interest to both clinicians and vision researchers for quite some time; the result is a variety of strongly held theories that tend to be contradictory. Some recent, excellently detailed reviews (Nelson, 1981; Jennings, 1985) have addressed various areas of research. However, anomalous correspondence transcends several research arenas; the object of this review is to correlate findings from several disparate disciplines into one parcel.

Binocular single vision normally is obtained on the horopter, a locus of points in space whose retinal images fall on pairs of corresponding points. A stimulus affecting one corresponding point will be localized in the same direction as if it had affected the other. The horopter, however, is a limited construct for describing the process of binocular, single vision; a disparity range can be added via sensory fusion. Moreover, since sensory fusion does not cover a large range either, suppression can be used as a further supplement to avoid physiological diplopia. This normal use of suppression can be enhanced by rivalrous conditions and has been postulated to play a role in strabismus (Halden, 1982).

Strabismus is said to be present when the line of sight of one eye fails to intersect the object of regard. When strabismus occurs in early childhood, certain adaptations can occur. Possible adaptations include amblyopia, suppression, and/or anomalous retinal correspondence. This paper will be limited to a discussion of anomalous retinal correspondence. However, it is clear that strict isolation of the possible adaptations is difficult.

Normal retinal correspondence (NRC) has been specified in a number of different ways. It has been defined as a physical point-to-point matching of the retinae of the two eyes

with fovea-to-fovea correspondence (Davson, 1972). Similarly, it has been defined as a correlation of retinal points with identical local signs or visual directions, with the foveal directions designating "straight-ahead" when at least the dominant eye is in the primary position (Moses, 1970). That is, an image falling on the right fovea will be perceived to be in the same direction as an image falling on the left fovea. This normal correspondence can occur independent of ocular alignment. Electrophysiologically NRC has been defined as the convergence of essentially monocular neurons upon a single binocular unit in striate cortex (Nelson, 1981).

Anomalous retinal correspondence (ARC) describes the condition in which a control mark seen only by the deviating eye of a squinter is perceived in the binocular visual field in a direction different than that to be expected on the basis of normal retinal correspondence (Moses, 1970). The spatial localization of the deviating eye appears to have partially or fully shifted, so as to counteract the effect of the ocular deviation. This shift in directional localization is termed the angle of anomaly. If the angle of anomaly equals the angle of squint, then the anomalous correspondence compensates exactly for the squint and is described as harmonious ARC (HAC). If the angle of anomaly is greater than zero, but less than the angle of squint, then it is described as unharmonious ARC (UHAC). If the angle of anomaly is in the opposite or noncompensating direction, the correspondence is described as paradoxical ARC (PAC) (Borish, 1970).

THREE COMMON TESTING TECHNIQUES

A variety of techniques, designed to evaluate retinal correspondence, have been devised. Each testing methodology appears to be influenced by the specific definition of correspondence accepted by the developer. Testing techniques,

in general, can be reviewed in Borish's *Clinical Refraction* (1970), from which three common testing methods are outlined here (Table 1).

The Hering or Bielschowsky (1937) after-image test is often used clinically because of its simplicity and apparent effectiveness in determining retinal correspondence. A bright lumi-line filament or flash attachment is used, with an opaque spot or band at the midpoint of the filament. The subject fixates the center point monocularly in order to achieve the after-image. This is commonly done with different orientations for each eye (note: For correct interpretation of this test normal or eccentric fixation must be determined prior to application.) If correspondence is normal the two after-images will form a cross; if correspondence is anomalous the two fixation points will be separated.

An amblyoscope can be used to determine the objective and subjective angles of deviation. By presenting differently oriented linear targets to each eye via separate arms of the amblyoscope, measurements can be made quickly and easily. Problems, involving either failure of image superimposition or suppression of one image, may occur and prevent the diagnosis of ARC.

The Bagolini striated glass test (Bagolini and Capobianco, 1965) makes use of plano lenses upon which striations 0.005 mm in width have been inscribed. The striations are uniform over each eye, but the axes of the striations are oriented 90 degrees away from each other. The viewing of a small fixation light creates the subjective observation of two blurred streaks with the rest of the visual scene appearing undistorted. If the two streaks cross at the fixated light, then the patient is exhibiting either NRC or ARC, depending on whether a squint is absent or present, respectively. Suppression can be a problem, but less of one than for the

amblyoscope. A small suppression zone, not noticed by the patient, can be missed if a critical examination is not performed. By avoiding dissociation, this test allows a determination of correspondence under normal visual conditions.

GENERALIZED THEORIES

The earliest theories considered ARC to be innate and immutable, reflecting a congenital anomaly that was untreatable (de la Hire, 1730; Muller, 1826). Later support for this view suggested that ARC was the underlying cause of squint in cases of large angle, nonaccommodative squints (Adler and Jackson, 1947). Failure of normal correspondence to develop postsurgically also has been blamed on a congenital ARC (Bedrossian, 1954).

Later theories held an alternate view that retinal correspondence changed as a sensory adaptation to the motor error or squint so that there was a shift in the visual direction of one eye relative to the other eye (Burian, 1947). The basic idea was that correspondence could be adaptively modified over some learning period in early life (Walls, 1951). Burian popularized the idea that ARC becomes more deeply ingrained over time and that this can be quantified or graded in terms of the nature of the stimulus conditions under which it persists. A common testing hierarchy has been suggested to test the "depth" of ARC adaptation: the Bagolini striated-glass test being proposed as the most sensitive means of eliciting ARC, the synoptophore or amblyoscope test purportedly exhibiting a mid-range sensitivity, and the after-image test being the least sensitive. Patients exhibiting ARC on the after-image test were taken to have a deeply embedded ARC of long standing (Mallett, 1970; Bagolini, 1976). However, the depth of anomaly theories have been questioned

Table 1
Three Common Testing Techniques

TEST	DEVICE	PROCEDURES	RESULTS
After-image test	Bright Lumi-line filament with an opaque band at the center.	Fixate horizontal filament with preferred eye. Then fixate vertical filament with nonpreferred eye.	NRC: a cross is seen. ARC: the two lines are apart.
Amblyoscope	Major Amblyoscope	Set arms to objective angle of squint. Present a vertical line to one eye, and a horizontal line to the other.	NRC: a cross is seen. ARC: the two lines are apart.
		Present lines, let patient set arms so that a cross is formed.	NRC: if the subjective angle equals the objective angle. ARC: subjective and objective angles are unequal.
Bagolini lens test	Striated lenses and a penlight.	Fixate a small penlight while looking thru the striated lenses with axes oriented OD-45; OS-135.	Squint present: ARC, if luminous streaks cross at fixation point. NRC, if they do not. Squint absent: NRC, if luminous streaks cross at fixation point. ARC if they do not.

(Flom and Kerr, 1967) based on measurement errors, unsteady eccentric fixation, and changes in the relative position of the eyes.

Another theory was the replacement theory of Verhoeff (1935, 1938); retinal correspondence was said not to be geared to subserve fusion, but merely to perceptually relate points on each retina on an alternating basis. This idea represented the first modern rivalry theory of vision. Brock (1941) essentially agreed with Verhoeff, suggesting that anomalous cases develop the ability to identify spatial localization separately with each eye. As an extension of Verhoeff's replacement theory Travers (1938, 1940) proposed that the diplopia of a squinter was relieved by the development of a suppression scotoma at the fovea, which sequentially spread across the eye until the development of an anomalous correspondence resulted. Since this was presumably a slow process, findings of unharmonious ARC, as well as harmonious ARC, easily could be explained. The implication was that the harmonious ARC was a completely adapted state, while the unharmonious ARC was a condition that was still in transition (Ronne and Rindziunski, 1953; Cashell and Durran, 1980). It was therefore assumed, similar to the Burian theory, that the longer standing and more constant the squint, the deeper established the ARC will become (Mallett, 1970; Bagolini, 1976).

An alternative adaptive theory has been proposed by Nelson (1981) suggesting a disparity "modal tuning" mechanism that responds globally to the most active disparity contour. As a result, there is a wide range of potentially corresponding states; specific conditions govern which correspondence state would be expressed at any one moment. An interpretation of Nelson's theory is similar to the hysteresis effect reported by Fender and Julesz (1967) with random dot stereograms. Once fusion is established, it can be maintained through an induced range of increasing disparity; once fusion is lost, the disparity must be decreased considerably before fusion can be reestablished. There may be, perhaps, a flexible and responsive stretching or shifting in correspondence that is capable of occurring only under specific conditions. Indeed, Campos (1982) has shown that strabismics have a wider Panum's area than normals. As are several other theories, this is consonant with harmonious ARC patients remaining harmonious despite the angle of squint changing from far to near, looking up and down (Mallett, 1967; Bagolini, 1976), or with and without spectacles (von Noorden, 1967). In support of the responsive correspondence shift theory is the finding of Maraini and Santori (1967) that ARC spontaneously may shift to NRC when the patient is forced to fixate with the normally deviating eye.

Nelson (1981) also proposed an alternative explanation for the varied test results that Burian ascribed to a deepened adaptation. Nelson suggested that the varied stimulus conditions differentially stimulate a global match response. The stimulus condition that best allows global matching to occur will be the most sensitive means of eliciting ARC; conversely, stimulus conditions that prevent global matching will fail to elicit ARC. Others have suggested the varied test results to

be totally artefactual because the different tests are evaluating different aspects of visual system performance (Duke-Elder, 1949).

Counter to all the adaptive or sensory theories, a motor theory has evolved suggesting an ARC that varies with the sensed position of the squinting eye. Briefly, eye position information could potentially change correspondence by influencing the correction from oculo-centric to ego-centric localization. In other words, the motor innervational pattern is proprioceptively registered so that the "turned eye" reports that what it sees is displaced with respect to the other eye (Duane, 1932). There is debate as to whether there is an adequate proprioceptive system within the extraocular musculature to provide such registration. Morgan (1961) considered eye movements and eye position to be registered because of innervation to the extraocular muscles; in effect an efference copy signal.

Boeder (1964, 1966) proposed a slightly different motor theory whereby a response shift always results when the eye is in a position different from the one called for by ocular innervation. Boeder feels that confirmation of his response shift hypothesis can be found in certain visual attributes of the strabismic patient: past pointing, monocular diplopia, and strabismic amblyopia.

Motor theories, in general, provide for an incorporation of an awareness by the visual system of some sort of kinesthetic signal, or an efference copy signal which allows a correspondence shift to take place. General motor theory evidence is the maintenance of correspondence when an eye is externally moved by tweezers grasped onto the conjunctiva when under local anesthesia; such an experiment acts to eliminate both proprioception and efference copy (Pasino and Maraini, 1965).

Both general theory types, adaptive and motor, have specific merit; clearly additional work is needed in order to reconcile the data. Initial attempts to "tease out" the determining factors have met with mixed results.

One means of differentiating the underlying issues is to determine whether there is a fusional movement of the eye or a change in correspondence when anomalous binocular vision is maintained under altered viewing conditions. Maraini and Pasino (1964) tried to separate the two possibilities. An after-image produced around the fovea of an ARC subject's deviating eye was located with respect to a fixation light viewed binocularly through Bagolini-stricted lenses. The introduction of six prism diopters base-out before the deviating eye of an ARC strabismic would have two possible effects if anomalous binocular single vision is maintained. If the after-image to fixation light separation remains constant, then a compensatory fusional movement has occurred. If the after-image to fixation light separation decreases by six prism diopters, then a compensatory shift of correspondence has occurred. Of eleven subjects evaluated, three showed a compensatory fusional movement, four showed a compensatory shift in correspondence, and the others showed a combination of the two. Other studies have shown similar conflicting results (Johnston, 1970; Kerr,

1980; Kenyon, Ciuffreda, and Stark, 1981). Since neither adaptive nor motor theories could be ruled out by this investigative means, it is unknown whether they apply to extremes of a single population distribution, or to separate physiological conditions. A similar parallel can be found elsewhere in the vision literature; anisometropic amblyopia and strabismic amblyopia have been suggested to represent separate conditions with overlapping symptoms.

All of the issues raised thus far have been based on results using subjective methodologies. Issue clarification might be obtained via objective evaluation techniques. Visual evoked potential data could prove to be useful since scalp topography of the VEP is known to reflect retinotopic mapping in the visual cortex (Jeffreys and Axford, 1972). Campos (1980) has detected binocular enhancement of pattern VEPs in ARC subjects, and also has routinely found binocular summation of VEPs in the same subjects (Campos and Chiesi, 1983; Chiesi et al, 1984). Yet, psychophysical investigations have failed to reveal any kind of summation in ARC subjects (Levi, 1987); very recent electrophysiological data, as well, have suggested that VEP summation is not representative of the activation of binocular cortical neurons (Shea, Aslin, and McCulloch, 1987). Additionally, McCormack (1987) has demonstrated that VEP scalp topographies do not differ for separate foveal stimulation of each eye in an ARC subject.

LABORATORY STUDIES

The field of electrophysiology also has attempted to analyze the processes governing the development of ARC. Animal experiments have suggested certain criteria are necessary for ARC development (Nelson, 1981): strabismus onset must be within the binocularity critical period, the strabismus must be small or gradually increasing, and cortical input competition must be balanced by having equal levels of monocular stimulation. The last point is perhaps the most important.

It has been shown that there is considerable overlap in ocular projections to the striate cortex that have withdrawn by termination of the binocularity critical period under normal conditions: induced strabismus has been shown to create an ocular dominance shift with the nondeviating eye gaining cortical connectivity at the expense of the deviating eye (Hubel, Wiesel, and LeVay, 1977). The presence of a strabismic condition, which meets the denoted criteria for ARC development, might interfere with the process of ocular dominance shifting, so that when the critical period is elapsed there are multiple, overlapping connections that are not present in strabismus with NRC.

Surgically induced strabismus in both cat and monkey has been shown to differentially damage the "X" visual subsystem (Brown and Salinger, 1975; Ikeda and Wright, 1976). Since this subsystem is characterized by exceptional spatial resolution and by fine disparity sensitivity (Poggio and Fischer, 1977), it is possible that strabismics exhibiting ARC are responding primarily via the "Y" visual subsystem. Since the "X" and "Y" subsystems have been postulated to provide the substrates for what has been termed "fine" and "coarse" stereopsis, respectively (Bishop and Henry, 1971), it is not

surprising that certain strabismics exhibiting ARC have a rudimentary stereopsis in the threshold range of about 100 sec. of arc and exhibit some coarse peripheral depth sensitivity (Helveston and von Noorden, 1967; Epstein and Tredici, 1973; Henson and Williams, 1980; Sireteanu, 1982).

If it is accepted that there might be multiple, overlapping ocular projections to the cortex, and if those inputs might be predominantly of the "Y-type" classification, then it would be reasonable to suggest that there is a dual neural alteration in strabismics with ARC. This dual alteration would exhibit itself as being able to provide only coarse, peripheral stereopsis; however, within this coarse framework the visual system may be able to shift correspondence tuning within the ranges of the broadly overlapping cortical projections, along the lines of Nelson's sensory fusion theory (1981). The resulting correspondence would only be "anomalous" with respect to the expected fusional limits demonstrated by "normals." Since "Y cell" input is provided both to the cortex and to the superior colliculus, it is possible that some sort of signal gating might be responsible for a sensory system and motor system match. In this manner both sensory adaptations and motor adaptations could work in tandem to provide for a shifting in correspondency. The concept, however, does not fully address motor theory issues, which are very important.

THERAPY

Theories regarding the treatment of ARC are as complicated and confusing as the theories regarding the underlying mechanism. The specific therapy techniques can include: the application of prisms, the prescribing of partially corrective lenses, the stimulation of stereopsis in a peripheral to central fashion, occlusion, bi-foveal stimulation (either in or out of instrumentation), the use of after-images, auditory biofeedback, as well as surgery. A large part of the literature stems from anecdotal clinical case reports.

Many of the treatment modes have approached therapy with respect to the motor theories (Boman and Kertesz, 1985; Kertesz and Kertesz, 1986). Vergence movements, either fusional (Kerr, 1968) or accommodative (Daum, 1982), may be registered by the visual system, and could thus be manipulated to stimulate ocular alignment. Sensory theory applications have also produced success: traditional amblyoscope therapy has been used to establish NRC superimposition (Griffin, 1976). Other treatment modes have sought to employ both motor and sensory aspects (Brock, 1941; Walraven, 1957; Ludlam, 1961; Wick, 1974 and 1975).

Treatment success is, however, another area that is open to debate. Reported cure rates have varied from 5% (Flom, 1969) to 85% (Pigassou-Albouy, 1973). Unfortunately, the definition of a cure varies from study to study. Other important factors that obscure the effect of treatment are: the method of ARC diagnosis, and the means of demonstrating NRC. Further difficulties consist of: a lack of clinical trials, possible experimenter bias, and poor experimental design. Future work will need to address these issues.

The implication is that the management of ARC is not a

simple manner. Careful diagnostic testing, under a variety of conditions, is necessary to correctly confirm the presence of ARC. Therapy requires the integration of both adaptive and motor techniques in order to maximize potential effectively.

SUMMARY

Theories of ARC fall into two broad categories: sensory adaptation, and motor response changes. Up to this point treatment modes have met with minimal overall success, largely as a result of the lack of any kind of consolidated model for ARC. Certainly, a major, organized, investigational effort is required if these issues are to be resolved.

REFERENCES

- Adler, F.H. and Jackson, F.E. (1947) Correlations between sensory and motor disturbances in convergent squint. *Arch. Ophthalmol.* 38:289-300.
- Begolius, B. and Capobianco, N. (1965) Subjective space in constant squint. *Am. J. Ophthalmol.* 59:430-442.
- Begolius, B. (1976) Part I. Sensory anomalies in strabismus (suppression, anomalous correspondence, amblyopia). *Doc. Ophthalmol.* 41:1-22. Part II. Sensory-motor anomalies in strabismus (anomalous movements). *Doc. Ophthalmol.* 41:23-41.
- Bedrossian, E.H. (1954) Anomalous retinal correspondence in alternating strabismus. *Arch. Ophthalmol.* 52:669-682.
- Bielachowski, A. (1937) Application of the after-image test in the investigation of squint. *Arch. Ophthalmol.* 17:3.
- Bishop, P.O. and Henry, G.H. (1971) Spatial vision. *Ann. Rev. Psychol.* 22:119-160.
- Boeder, P. (1964) Anomalous retinal correspondence revised. *Am. J. Ophthalmol.* 58:366-373.
- Boeder, P. (1966) Single binocular vision in strabismus. *Am. J. Ophthalmol.* 61:78-86.
- Bornes, D.K. and Kertesz, A.E. (1985) Fusional responses of strabismic subjects to foveal and extra-foveal stimulation. *Invest. Ophthalmol. Vis. Sci.* 26:1731-1739.
- Bornes, I.M. (1970) *Clinical Refraction*, 3rd ed. pp 1233-1240. Professional Press, Inc., Chicago.
- Brock, F.W. (1941) Conditioning the squinter to normal visual habits. *Optom. Weekly* 32:793-801; 819-824.
- Brown, D.L. and Selinger, W.L. (1975) Loss of X-cells in lateral geniculate nucleus with monocular paralysis: neural plasticity in the adult cat. *Science* 189:1011-1012.
- Burian, H.M. (1947) Sensory retinal relationships in concomitant strabismus. *Arch. Ophthalmol.* 37:336-368; 504-533; 618-648.
- Campos, E.C. (1980) Anomalous retinal correspondence: monocular and binocular visual evoked responses. *Arch. Ophthalmol.* 98:299-302.
- Campos, E.C. (1982) Binocularity in constant strabismus: binocular visual field studies. *Doc. Ophthalmol.* 53:249-281.
- Campos, E.C. and Chien, C. (1983) Binocularity in constant strabismus. *Doc. Ophthalmol.* 55:277-293.
- Cashell, G.T.W. and Durran, I.M. (1980) In *Handbook of Orthoptic Principles*, 4th ed. Churchill Livingstone, Edinburgh.
- Chien, C., Sargente, A.D., and Bolzano, R. (1984) Binocular visual perception in strabismus studied by means of visual evoked responses. *Doc. Ophthalmol.* 58:51-56.
- de la Hère, (1730) Quoted in Halden U. (1952). Fusional phenomena in anomalous correspondence. *Acta Ophthalmol. Suppl.* 37:1-93.
- Dennis, K.M. (1982) Covariation in anomalous correspondence with accommodative vergence. *Am. J. Optom. Physiol. Opt.* 59:146-151.
- Davson, H. (1972) *The Physiology of the Eye*. Academic Press, New York, p. 447.
- Dennis, A. (1932) Diplopia and other disorders of binocular projection. *Arch. Ophthalmol.* 7:187-210.
- Duke Elder, W. (1949) *Textbook of Ophthalmology*. C.V. Mosby Co., St. Louis.
- Eppstein, D.L. and Tradici, T.J. (1973) Microtropia in flying personnel. *Am. J. Ophthalmol.* 76: 832-841.
- Fender, D. and Julez, B. (1967) Extension of Panum's fusional area in binocularly stabilized vision. *J. Opt. Soc. Am.* 57:819-830.
- Flom, M.C. (1969) In Hirsch, M.J. et al (eds). *Vision of Children*. Chilton Book Co., Philadelphia.
- Flom, M.C. and Kerr, K.E. (1967) Determination of retinal correspondence. *Arch. Ophthalmol.* 77:200-213.
- Griffith, J.R. (1976) *Binocular Anomalies: Procedures for Vision Therapy*. Chicago Professional Press, Inc. pp. 242-245.
- Halden, U. (1962) Suppression accommodation in concomitant strabismus with harmonious anomalous correspondence. *Acta Ophthalmol.* 60:828-834.
- Helveston, E.M. and Von Noorden, G.K. (1967) Microtropia: a newly defined entity. *Arch. Ophthalmol.* 78:272-281.
- Hesslen, D.B. and Williams, D.E. (1980) Depth perception in strabismus. *Brit. J. Ophthalmol.* 64:349-353.
- Hubel, D.H., Wiesel, T.N., and LeVay, S. (1977) Plasticity of ocular dominance columns in monkey striate cortex. *Phil. Trans. Roy. Soc. London B.* 278:377-409.
- Ikeda, H. and Wright, M.J. (1976) Properties of LGN cells in kittens reared with convergent squint: a neurophysiological demonstration of amblyopia. *Exp. Brain Res.* 25:63-77.
- Jeffreys, D.A. and Axford, J.G. (1972) Source localizations of pattern-specific components of human visual evoked potentials.
- I. Component of striate cortical origin. *Exp. Brain Res.* 16: 1-21.
 - II. Component of extrastriate cortical origin. *Exp. Brain Res.* 16: 22-40.
- Jennings, J.A.M. (1985) Anomalous retinal correspondence - a review. *Ophthalmol. Physiol. Opt.* 5:357-368.
- Johnston, A.W. (1970) An analysis of clinical test results for anomalous correspondence. *Aust. J. Optom.* 53:38-56.
- Kanyon, R.V., Ciuffreda, K.J., and Stark, L. (1981) Dynamic vergence eye movements in strabismus and amblyopia: asymmetric vergence. *Brit. J. Ophthalmol.* 65:167-176.
- Kerr, K.E. (1968) Instability of anomalous retinal correspondence. *J. Am. Optom. Assoc.* 39:1107-1108.
- Kerr, K. (1980) Accommodative and fusional vergence in anomalous correspondence. *Am. J. Optom. Physiol. Opt.* 57: 676-680.
- Kertesz, A.E. and Kertesz, J. (1986) Wide-field fusional stimulation in strabismus. *Am. J. Optom. Physiol. Opt.* 63: 217-222.
- Levi, D. (1987) Personal report.
- Ludman, W.M. (1961) Orthoptic treatment of strabismus. *Am. J. Optom. Arch. Acad. Optom.* 38:369-388.
- Mallett, R.F.J. (1967) ARC and Bagolini striated lenses. *Ophthalmol. Optician* 7:164-166.
- Mallett, R.F.J. (1970) Anomalous correspondence - the new outlook. *Ophthalmol. Optician* 10:606-608; 621-624.
- Martini, G. and Finino, L. (1964) Variations in the angle of anomaly and fusional movements in cases of small angle convergent strabismus with harmonious anomalous retinal correspondence. *Brit. J. Ophthalmol.* 48:439-443.
- Marsden, G. and Satter, M. (1967) Anomalous retinal correspondence and monocular squint. *Ophthalmologica* 153:179-183.
- McCormick, G. (1988) Normal retinotopic mapping in human strabismus with ARC. In review.
- Morgan, M.W. (1961) Anomalous correspondence interpreted as a motor phenomenon. *Am. J. Optom. Arch. Acad. Optom.* 38:131-148.
- Moore, R.A. (1970) *Adler's Physiology of the Eye*. The C.V. Mosby Co., Saint Louis, pp. 138 and 668.
- Müller, H. (1826) In Donders, F.C. (1864) On the anomalies of accommodation and refraction of the eye. pp. 250-251. The New Sydenham Society, London.
- Nelson, J.L. (1981) A neurophysiological model for anomalous correspondence based on mechanisms of sensory fusion. *Doc. Ophthalmol.* 51:3-99.
- Pasino, L. and Marsden, G. (1965) Unpublished data referred to in the previous references.
- Pigganov-Albeoy, R. (1973) The use of over-correcting prisms in the treatment of strabismic patients without amblyopia. *Albrecht von Graefes Arch. Klin. Ophthalmol.* 186:209-226.
- Poggio, G.F. and Fischer, B. (1977) Binocular interaction and depth sensitivity in striate and pre-striate cortex of behaving rhesus monkey. *J. Neurophysiol.* 40:1392-1405.
- Ronne, G. and Rindzinski, E. (1953) The diagnosis and clinical classification of anomalous correspondence. *Acta Ophthalmol.* 31:321-345.
- Shaw, S.L., Astin, R.N., and McCulloch, D. (1987) Binocular VEP summation in infants and adults with abnormal binocular histories. *Invest. Ophthalmol. Vis. Sci.* 28:356-365.
- Sireteanu, R. (1982) Binocular vision in strabismic humans with alternating fixation. *Vision Res.* 22: 889-896.
- Travers, T.A.B. (1938) Suppression of vision in squint and its association with retinal correspondence and amblyopia. *Brit. J. Ophthalmol.* 22:577-604.
- Travers, T.A.B. (1940) The origin of abnormal retinal correspondence. *Brit. J. Ophthalmol.* 24:58-64.
- Verhoeff, F.H. (1935) A new theory of binocular vision. *Arch. Ophthalmol.* 13:151-175.
- Verhoeff, F.H. (1938) Anomalous projection and other visual phenomena associated with strabismus. *Arch. Ophthalmol.* 19:663-699.
- von Noorden, G.K. (1967) Discussion of Bagolini. *Doc Ophthalmol.* 23:388.
- Walls, G.L. (1951) The problem of visual direction.
- Part I. The history to 1900. *Am. J. Optom.* 28:55-83.
 - Part II. The tangible basis for nearvision. *Am. J. Optom.* 28:115-146.
 - Part III. Experimental attacks and their results. *Am. J. Optom.* 28:173-212.
- Walraven, F.C. (1957) A discussion of anomalous retinal correspondence. *Am. Orthop. J.* 7:162-175.
- Wick, B. (1974) Visual therapy for constant exotropia with anomalous retinal correspondence - a case report. *Am. J. Optom. Physiol. Opt.* 51:1005-1008.
- Wick, B. (1975) "Forced elimination" of anomalous retinal correspondence in constant exotropia - a case report. *Am. J. Optom. Physiol. Opt.* 52:58-62.

AUTHOR'S ADDRESS:

P.O. Box 577
Fort Rucker, AL 36362-5292

Initial distribution

Commander
U.S. Army Natick Research
and Development Center
ATTN: Documents Librarian
Natick, MA 01760

Naval Submarine Medical
Research Laboratory
Medical Library, Naval Sub Base
Box 900
Groton, CT 05340

Commander/Director
U.S. Army Combat Surveillance
& Target Acquisition Lab
ATTN: DELCS-D
Fort Monmouth, NJ 07703-5304

Commander
10th Medical Laboratory
ATTN: Audiologist
APO NEW YORK 09180

Commander
Naval Air Development Center
Biophysics Lab
ATTN: G. Kydd
Code 60B1
Warminster, PA 18974

Naval Air Development Center
Technical Information Division
Technical Support Detachment
Warminster, PA 18974 -

Dr. E. Hendler
Human Factors Applications, Inc.
295 West Street Road
Warminster, PA 18974

Under Secretary of Defense
for Research and Engineering
ATTN: Military Assistant
for Medical and Life Sciences
Washington, DC 20301

Commander
U.S. Army Research Institute
of Environmental Medicine
Natick, MA 01760

U.S. Army Avionics Research
and Development Activity
ATTN: SAVAA-P-TP
Fort Monmouth, NJ 07703-5401

U.S. Army Research and Development
Support Activity
Fort Monmouth, NJ 07703

Chief, Benet Weapons Laboratory
LCWSL, USA ARRADCOM
ATTN: DRDAR-LCB-TL
Watervliet Arsenal, NY 12189

Commander
Man-Machine Integration System
Code 602
Naval Air Development Center
Warminster, PA 18974

Commander
Naval Air Development Center
ATTN: Code 6021 (Mr. Brindle)
Warminster, PA 18974

Commanding Officer
Naval Medical Research
and Development Command
National Naval Medical Center
Bethesda, MD 20014

Director
Army Audiology and Speech Center
Walter Reed Army Medical Center
Washington, DC 20307-5001

COL Carl F. Tyner, MC
Walter Reed Army Institute
of Research
Washington, DC 20307-5100

HQ DA (DASG-PSP-0)
5109 Leesburg Pike
Falls Church, VA 22041-3258

Naval Research
Laboratory Library
Code 1433
Washington, DC 20375

Harry Diamond Laboratories
ATTN: Technical Information Branch
2800 Powder Mill Road
Adelphi, MD 20783-1197

U.S. Army Materiel Systems
Analysis Agency
ATTN: Reports Processing
Aberdeen Proving Ground
MD 21005-5017

U.S. Army Ordnance Center
and School Library
Building 3071
Aberdeen Proving Ground,
MD 21005-5201

U.S. Army Environmental Hygiene
Agency
Building E2100
Aberdeen Proving Ground,
MD 21010

Technical Library
Chemical Research
and Development Center
Aberdeen Proving Ground,
MD 21010-5423

Commander
U.S. Army Institute
of Dental Research
Walter Reed Army Medical Center
Washington, DC 20307-5300

Naval Air Systems Command
Technical Air Library 950D
Rm 278, Jefferson Plaza II
Department of the Navy
Washington, DC 20361

Naval Research Laboratory Library
Shock and Vibration Information Center,
Code 5804
Washington, DC 20375

Director
U.S. Army Human Engineering Laboratory
ATTN: Technical Library
Aberdeen Proving Ground,
MD 21005-5001

Commander
U.S. Army Test
and Evaluation Command
ATTN: AMSTE-AD-H
Aberdeen Proving Ground,
MD 21005-5055

Director
U.S. Army Ballistic
Research Laboratory
ATTN: DRXBR-OD-ST Tech Reports
Aberdeen Proving Ground,
MD 21005-5066

Commander
U.S. Army Medical Research
Institute of Chemical Defense
ATTN: SGRD-UV-AO
Aberdeen Proving Ground,
MD 21010-5425

Commander
U.S. Army Medical Research
and Development Command
ATTN: SGRD-RMS (Ms. Madigan)
Fort Detrick, Frederick, MD 21701

Commander
U.S. Army Medical Research
Institute of Infectious Diseases
Fort Detrick, Frederick,
MD 21701

Director, Biological
Sciences Division
Office of Naval Research
600 North Quincy Street
Arlington, VA 22217

Commander
U.S. Army Materiel Command
ATTN: AMCDE-S (CPT Broadwater)
5001 Eisenhower Avenue
Alexandria, VA 22333

Commandant
U.S. Army Aviation
Logistics School
ATTN: ATSQ-TDN
Fort Eustis, VA 23604

U.S. Army Training
and Doctrine Command
ATTN: ATCD-ZX
Fort Monroe, VA 23651

Structures Laboratory Library
USARTL-AVSCOM
NASA Langley Research Center
Mail Stop 266
Hampton, VA 23665

Naval Aerospace Medical
Institute Library
Bldg 1953, Code 102
Pensacola, FL 32508

Command Surgeon
U.S. Central Command
MacDill Air Force Base
FL 33608

Air University Library
(AUL/LSE)
Maxwell AFB, AL 36112

Commander
U.S. Army Biomedical Research
and Development Laboratory
ATTN: SGRD-UBZ-I
Fort Detrick, Frederick,
MD 21701

Defense Technical
Information Center
Cameron Station
Alexandria, VA 22313

U.S. Army Foreign Science
and Technology Center
ATTN: MTZ
220 7th Street, NE
Charlottesville, VA 22901-5396

Director,
Applied Technology Laboratory
USARTL-AVSCOM
ATTN: Library, Building 401
Fort Eustis, VA 23604

U.S. Army Training
and Doctrine Command
ATTN: Surgeon
Fort Monroe, VA 23651-5000

Aviation Medicine Clinic
TMC #22, SAAF
Fort Bragg, NC 28305

U.S. Air Force Armament
Development and Test Center
Eglin Air Force Base, FL 32542

U.S. Army Missile Command
Redstone Scientific
Information Center
ATTN: Documents Section
Redstone Arsenal, AL 35898-5241

U.S. Army Research and Technology
Laboratories (AVSCOM)
Propulsion Laboratory MS 302-2
NASA Lewis Research Center
Cleveland, OH 44135

AFAMRL/HEX
Wright-Patterson AFB, OH 45433

University of Michigan
NASA Center of Excellence
in Man-Systems Research
ATTN: R. G. Snyder, Director
Ann Arbor, MI 48109

John A. Dellinger,
Southwest Research Institute
P. O. Box 28510
San Antonio, TX 78284

Project Officer
Aviation Life Support Equipment
ATTN: AMCPO-ALSE
4300 Goodfellow Blvd.
St. Louis, MO 63120-1798

Commander
U.S. Army Aviation
Systems Command
ATTN: DRSAB-ED
4300 Goodfellow Blvd
St. Louis, MO 63120

Commanding Officer
Naval Biodynamics Laboratory
P.O. Box 24907
New Orleans, LA 70189

U.S. Army Field Artillery School
ATTN: Library
Snow Hall, Room 14
Fort Sill, OK 73503

Commander
U.S. Army Health Services Command
ATTN: HSOP-SO
Fort Sam Houston, TX 78234-6000

U.S. Air Force Institute
of Technology (AFIT/LDEE)
Building 640, Area B
Wright-Patterson AFB, OH 45433

Henry L. Taylor
Director, Institute of Aviation
University of Illinois-
Willard Airport
Savoy, IL 61874

Commander
U.S. Army Aviation
Systems Command
ATTN: DRSAB-WS
4300 Goodfellow Blvd
St. Louis, MO 63120-1798

Commander
U.S. Army Aviation
Systems Command
ATTN: SGRD-UAX-AL (MAJ Lacy)
4300 Goodfellow Blvd., Bldg 105
St. Louis, MO 63120

U.S. Army Aviation
Systems Command
Library and Information
Center Branch
ATTN: DRSAB-DIL
4300 Goodfellow Blvd
St. Louis, MO 63120

Federal Aviation Administration
Civil Aeromedical Institute
CAMI Library AAC 64D1
P.O. Box 25082
Oklahoma City, OK 73125

Commander
U.S. Army Academy
of Health Sciences
ATTN: Library
Fort Sam Houston, TX 78234

Commander
U.S. Army Institute
of Surgical Research
ATTN: SGRD-USM (Jan Duke)
Fort Sam Houston, TX 78234-6200

Director of Professional Services
AFMSC/GSP
Brooks Air Force Base, TX 78235

U.S. Army Dugway Proving Ground
Technical Library
3Bldg 5330
Dugway, UT 84022

U.S. Army Yuma Proving Ground
Technical Library
Yuma, AZ 85364

AFFTC Technical Library
6520 TESTG/ENXL
Edwards Air Force Base,
CAL 93523-5000

Commander
Code 3431
Naval Weapons Center
China Lake, CA 93555

Aeromechanics Laboratory
U.S. Army Research
and Technical Labs
Ames Research Center,
M/S 215-1
Moffett Field, CA 94035

Sixth U.S. Army
ATTN: SMA
Presidio of San Francisco,
CA 94129

Commander
U.S. Army Aeromedical Center
Fort Rucker, AL 36362

Directorate
of Combat Developments
Bldg 507
Fort Rucker, AL 36362

U.S. Air Force School
of Aerospace Medicine
Strughold Aeromedical Library
Documents Section, USAFSAM/TSK-4
Brooks Air Force Base, TX 78235

Dr. Diane Damos
Department of Human Factors
ISSM, USC
Los Angeles, CA 90089-0021

U.S. Army White Sands
Missile Range
Technical Library Division
White Sands Missile Range,
NM 88002

U.S. Army Aviation Engineering
Flight Activity
ATTN: SAVTE-M (Tech Lib)
Stop 217
Edwards Air Force Base,
CA 93523-5000

U.S. Army Combat Developments
Experimental Center
Technical Information Center
Bldg 2925
Fort Ord, CA 93941-5000

Commander
Letterman Army Institute
of Research
ATTN: Medical Research Library
Presidio of San Francisco,
CA 94129

Director
Naval Biosciences Laboratory
Naval Supply Center, Bldg 844
Oakland, CA 94625

Commander
U.S. Army Medical Materiel
Development Activity
Fort Detrick,
Frederick, MD 21701-5009

Directorate
of Training Development
Bldg 502
Fort Rucker, AL 36362

Chief
Army Research Institute
Field Unit
Fort Rucker, AL 36362

Commander
U.S. Army Safety Center
Fort Rucker, AL 36362

U.S. Army Aircraft Development
Test Activity
ATTN: STEBG-MP-QA
Cairns AAF
Fort Rucker, AL 36362

Naval Aerospace Medical
Institute Library
Building 1953, Code 102
Pensacola, FL 32508

Commander
U.S. Army Medical Research
and Development Command
ATTN: SGRD-PLC (COL Sedge)
Fort Detrick, Frederick
MD 21701

Chief
Human Engineering Laboratory
Field Unit
Fort Rucker, AL 36362

Commander
U.S. Army Aviation Center
and Fort Rucker
ATTN: ATZQ-T-ATL
Fort Rucker, AL 36362

President
U.S. Army Aviation Board
Cairns AAF
Fort Rucker, AL 36362

Commanding Officer
Harry G. Armstrong Aerospace
Medical Research Lab
Wright-Patterson
Air Force Base, OH 45433